



OPEN ACCESS

Effects of city-wide 20 mph (30km/hour) speed limits on road injuries in Bristol, UK

Anna Bornioli ,¹ Isabelle Bray,¹ Paul Pilkington,¹ John Parkin²

¹Centre for Public Health and Wellbeing, Department of Health and Social Sciences, University of the West of England Bristol, Bristol, UK

²Centre for Transport and Society, Department of Geography and Environmental Management, University of the West of England Bristol, Bristol, UK

Correspondence to

Dr Anna Bornioli, Health and Social Sciences, University of the West of England Bristol, Bristol BS16 1QY, UK; anna.bornioli@uwe.ac.uk

Received 14 May 2019

Revised 25 June 2019

Accepted 26 June 2019

Published Online First

13 July 2019

ABSTRACT

Twenty miles per hour (32 km/hour) or 30 km/hour speed limits represent a potential strategy to reduce urban road injuries and are becoming increasingly widespread. However, no study has conducted a robust evaluation of the effects of city-wide 20mph speed limits on road injuries. This study reports the effects of such an intervention, based on a natural experiment that took place in Bristol, UK. Based on a stepped-wedge design using count data, negative binomial regressions showed that between 2008 and 2016, the 20 mph speed limit intervention was associated with a city-level reduction of fatal injuries of around 63% (95% CI 2% to 86%), controlling for trends over time and areas. There was also a general trend of reduction of the total number of injuries at city level and in 20 mph roads. These findings highlight the potential benefits of city-wide 20 mph speed limits. We hypothesise that this city-wide approach may encourage a general behaviour change in drivers that, in turn, may contribute to reducing injuries across the city.

INTRODUCTION

Twenty miles per hour (32 km/hour) or 30 km/hour speed limits represent a potential strategy to reduce urban road injuries and are becoming increasingly widespread in Western countries.^{1–3} This is because reducing road injuries is a priority for public health; as such, injuries are among the leading causes of death and disability worldwide, with more than one million individuals dying each year on the road.⁴ Previous research has shown that motorised vehicle speed is positively associated with risk of road traffic collisions^{5,6}; therefore, reducing speed should have an effect on reducing injuries.⁴ Twenty miles per hour limits are sign-only interventions without engineering measures, such as speed bumps, as opposed to 20mph speed zones, which also include speed-reduction engineering measures to slow traffic. According to the WHO, 20mph limits are recommended for residential areas with high pedestrian activity.⁵

In 2009, a study found that 20mph zones in London were associated with a 41.9% reduction in road casualties.⁷ However, despite 20mph speed limits (ie, without engineering measures) becoming increasingly popular, little research has been conducted to test the effects of 20mph limits on road injuries. A recent UK government study concluded that there is not enough evidence to suggest that 20mph speed limits are associated with decreases in injuries in residential areas.⁸ However, no study has conducted a robust evaluation of the

effects of a city-wide 20mph speed limit intervention, defined as a majority of roads across an urban area being set at 20mph, on road injuries. The implementation in Bristol, UK, was conceived in principle as being comprehensively city-wide, with exceptions for a few roads that remained at 30 mph. This paper reports findings from the first stepped-wedge design road injury evaluation of such an intervention. The study has the strengths of being based on a natural experiment and of being the first study to evaluate the effects of such a high-coverage policy on injuries with a robust methodology.

METHODS

The Bristol 20mph speed limit policy was implemented between 2010 and 2015.⁹ The 20mph limit scheme was introduced in phases corresponding to seven areas (for details on the intervention, refer to Bornioli *et al*¹⁰), enabling a stepped-wedge design analysis. This is a pragmatic design suited to interventions that are implemented in steps across several clusters (in this case, geographical areas). Such a design allows robust evaluation of the effects of natural experiments where interventions are implemented following political or logistic constraints.^{11,12}

The current injury analysis was based on Police STATS19 data from 2008 to 2016. STATS19 records the date, location, and number and type of injury (slight, serious or fatal). An injury is defined as serious if the person is detained in the hospital as an inpatient or has any of the following injuries: fractures, concussion, internal injuries, crushing, non-friction burns, severe cuts and lacerations, or severe general shock requiring medical treatment. An injury is classified as fatal if the person dies within 30 days of the collision.

In the injury dataset, the Bristol City Council coded each injury with its area code, severity, the intervention period (preintervention or postintervention), the speed limit of the road (20 or 30mph) and sociodemographics of the injured people. The research team further classified each collision by year and calculated exposure times for each area in each year (eg, the number of months included in the study either preintervention or postintervention in each area; table 1). Figure 1 illustrates the stepped-wedge study design.

Statistical methods

The analysis compared injury counts before and after the introduction of the 20mph speed limit while controlling for year and areas. A stepped-wedge design enables a robust evaluation based on a natural experiment, allowing for the rolling nature of the intervention, implemented in several



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Bornioli A, Bray I, Pilkington P, *et al.* *Inj Prev* 2020;**26**:85–88.

Table 1 Number of injuries and equivalent annual rates of injury for all roads

Area	Injury type	Months, rounded (n)		Injuries (n)		Equivalent annual rate of injuries	
		Pre	Post	Pre	Post	Pre	Post
1	All severity	34	74	601	1143	212.1	185.4
	Fatal	34	74	2	2	0.7	0.3
	Serious	34	74	60	116	21.2	18.8
	Slight	34	74	539	1025	190.2	166.2
2	All severity	72	36	2092	862	348.7	287.3
	Fatal	72	36	15	4	2.5	1.3
	Serious	72	36	228	105	38.0	35.0
	Slight	72	36	1849	753	308.2	251.0
3	All severity	78	30	1208	462	185.8	184.8
	Fatal	78	30	12	2	1.8	0.8
	Serious	78	30	138	65	21.2	26.0
	Slight	78	30	1058	395	162.8	158.0
4	All severity	81	27	939	243	139.1	108.0
	Fatal	81	27	19	3	2.8	1.3
	Serious	81	27	109	33	16.1	14.7
	Slight	81	27	811	207	120.1	92.0
5	All severity	87	21	1284	260	177.1	148.6
	Fatal	87	21	19	0	2.6	0.0
	Serious	87	21	116	28	16.0	16.0
	Slight	87	21	1149	232	158.5	132.6
6	All severity	89	19	631	107	85.1	67.6
	Fatal	89	19	4	1	0.5	0.6
	Serious	89	19	73	15	9.8	9.5
	Slight	89	19	554	91	74.7	57.5
7	All severity	93	15	852	133	109.9	106.4
	Fatal	93	15	9	0	1.2	0.0
	Serious	93	15	85	11	11.0	8.8
	Slight	93	15	758	122	97.8	97.6

steps.^{11 12} In addition, controlling for calendar year (as a continuous variable) allows for changes in injury rates over time that were not associated with the intervention (eg, national trends). Two analyses were undertaken: (1) at city level (including roads with postintervention limits of 20 and 30mph) and (2) on 20mph roads only. This is because previous analyses^{2 12} found that average speeds in Bristol decreased following the intervention from 27.1 to 23.7 mph, with postintervention mean speeds of 22.5 mph in 20mph roads and of 25.9 mph in 30mph roads. Analyses of injuries, rather than collisions, was undertaken because this is the primary outcome of public health interest.

Count data are often analysed using Poisson regression. However, the goodness-of-fit test in Stata 14 (`estat gof`) showed overdispersion. Therefore, negative binomial regression was used instead (`nbreg` command). Adjusted incident rate ratios (IRRs) were estimated to compare injuries preintervention

and postintervention. Analyses were run separately for each injury type (slight, serious and fatal). In 2016, a revision to the injury reporting regime called CRASH was introduced. This has resulted in an increase in the proportion of non-fatal injuries categorised as 'serious'.⁸ For this reason, sensitivity analyses were performed by re-running the analyses for serious and slight injuries excluding observations from year 2016. Results did not vary, but these are available on request.

Sites for traffic safety interventions are sometimes selected based on relative performance, with the worst performing sites being selected for treatment. If sites displayed high preintervention levels of collisions and injuries because of randomness, then over time, we would expect these sites to revert to a lower mean value.¹³ However, regression to the mean was not considered an issue in this study, as the 20 mph limits were introduced across the majority of the city's roads, rather than focused on known collision sites.

RESULTS

Descriptive analyses showed that annual postintervention rates of fatal, serious and slight injuries in each Bristol area are generally lower than the respective injury rate before the limits were introduced (table 1). Given that areas had different implementation dates, it was not possible to run descriptive analyses on city-level injury rates.

Inferential analysis

Table 2 summarises the estimated effects of the 20 mph limit on injuries. This shows a general trend of injury reductions


**Figure 1** Stepped-wedge study design.

Table 2 IRRs comparing injuries preintervention and postintervention, by injury type, for all roads and for 20 mph roads only

	Injury severity	IRR (95% CI)*	P value
All roads (20 and 30 mph roads)	Fatal	0.37 (0.14 to 0.98)	0.04
	Serious	1.02 (0.77 to 1.34)	0.86
	Slight	0.94 (0.84 to 1.06)	0.39
20 mph roads only	Fatal	0.65 (0.15 to 2.83)	0.57
	Serious	0.84 (0.60 to 1.17)	0.32
	Slight	0.91 (0.78 to 1.07)	0.26

*Controlling for calendar year (continuous variable) and areas (categorical variable).

IRR, incidence rate ratio.

following the intervention, both city-wide and in 20 mph roads only, and strong evidence of a city-wide reduction of fatal injuries of around 63%, though this estimate was imprecise (95% CI 2% to 86%). No temporal trend was detected.

DISCUSSION

The analysis highlights a general reduction in injuries and suggests evidence of a city-level reduction in fatalities of 63%. These findings are in line with the reductions of injuries in Brighton City Centre, where a similar area-wide (but not city-wide) scheme has been implemented.⁸ Interestingly, the reductions in fatalities in Bristol are larger than the 46% reduction in London associated with the introduction of traffic-calmed 20 mph zones, as identified by Grundy *et al.*⁷ The city-level reduction of fatal injuries identified in our study should also be set against national trends, which show that the number of deaths on built-up roads has increased from the 2010–2014 annual average of 585 to a 2017 figure of 607 deaths.¹⁴ Therefore, this finding goes some way to supporting city-wide 20 mph speed limits as an intervention to reduce road injuries.

The fact that such a solid reduction was observed on city level, rather than on 20 mph roads only, might be explained by the fact that average speeds decreased overall in Bristol following the intervention, and not only in 20 mph roads.^{2 12} Preintervention average speed in 30 mph roads was relatively high, and the reduction in speed may have contributed to reducing fatal injuries in these roads. This is similar to findings from Brighton, where reductions in collisions were observed especially in major strategic roads.⁸ This also suggests that the reduction in injuries is due to a general change in driving behaviour related to the city-wide nature of the 20 mph speed limit, with slower driving speeds also in 30 mph roads, where speed tends to be higher than in 20 mph roads.

To our knowledge, the current study is the first study examining the effects of a city-wide 20 mph speed limit. The study is also the first to apply a stepped-wedge design with injury count data for a 20 mph speed limit intervention, based on a natural experiment. While a recent national report has found that there is insufficient evidence to conclude that 20 mph speed limits implemented in specific city areas can reduce injuries, the current findings suggest there are benefits of city-wide 20 mph interventions. Therefore, city-wide measures, as opposed to more geographically restricted 20 mph speed limits, such as those targeting areas of perceived higher risk, outside schools for example, may help to reduce road injuries.

Limitations

A limitation of the analysis is the potential inaccuracy of STATS19 data, which are known to under-report injuries. A

second limitation is the relatively short time frame of the study, with 9 years of data available. However, time frames are similar to those of Department of Transport (DfT)⁸; more robust evidence will be available in the long term. Third, there was no randomisation in the design, but this is a common feature in natural experiments. In addition, the wide 95% CIs might be due to the relatively small number of injuries observed, particularly fatal injuries. Finally, although we controlled for linear trends in casualties over time, we could not include a comparator area controlling for other factors (eg, weather and changes in emergency care) as in DfT⁸ because the intervention was applied to the whole of the study. We recognise that the strength of the statistical analysis could have been improved by the inclusion of a comparator city without city-wide 20 mph limits.

CONCLUSIONS

The paper reported findings from the first stepped-wedge design evaluation of the effects of a city-wide 20 mph speed limit intervention on road injuries. Findings have identified a general trend of reductions in road injuries and strong evidence of a reduction of city-level fatal injuries following the 20 mph speed limit intervention. This suggests that city-wide 20 mph speed limits could be an effective strategy for reducing injuries, as it encourages safer driving behaviour that is not restricted to 20 mph roads. The policy recommendation that would follow from this is to implement city-wide 20 mph limits, rather than 20 mph limits in restricted areas of a city. Further research is needed and changes in injury rates should be monitored over longer time frames.

What is already known on the subject

- ▶ Motorised vehicle speed is positively associated with risk of road traffic injuries.
- ▶ Twenty miles per hour (30 km/hour) limits as a road safety tool to reduce casualties are increasingly popular.
- ▶ There is no city-wide robust evaluation of the effects of city-wide 20 mph speed limits on road injuries.

What this study adds

- ▶ This is the first study to apply a stepped-wedge design with count data to a city-wide 20 mph limit case study.
- ▶ The study identifies a general trend of injury reductions and strong evidence of reduction of fatal injuries following a city-wide 20 mph speed limit intervention.

Funding This study was supported by Bristol City Council.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Anna Bornioli <http://orcid.org/0000-0001-9452-2907>

REFERENCES

- 1 Hu W, Cicchino JB. Lowering the speed limit from 30 mph to 25 mph in Boston: effects on vehicle speeds. *Inj Prev* 2019. doi: 10.1136/injuryprev-2018-043025.
- 2 Pilkington P, Bornioli A, Bray I, *et al*. The Bristol twenty miles per hour limit evaluation (BRITE) study. *University of the West of England, Bristol* 2018.
- 3 Fridman L, Rothman L, Hagel B, *et al*. The effect of lowering residential speed limits to 30 km/h on child pedestrian injuries in Toronto, Canada. *Injury Prevention* 2018;24.
- 4 World Health Organisation. Global status report on road safety 2018 World Health Organisation; 2018.
- 5 Elvik R. *Speed and road safety: synthesis of evidence from evaluation studies*. Journal of the Transportation Research Board: Transportation Research Record, 2005: 1908. 59–69.
- 6 Rosén E, Stigson H, Sander U. Literature review of pedestrian fatality risk as a function of car impact speed. *Accid Anal Prev* 2011;43:25–33.
- 7 Grundy C, Steinbach R, Edwards P, *et al*. Effect of 20 mph traffic speed zones on road injuries in London, 1986–2006: controlled interrupted time series analysis. *BMJ* 2009;339.
- 8 Department of Transport. 20mph research study report: process and impact evaluation technical report; 2018.
- 9 Bristol City Council. 20mph speed limit pilot areas: monitoring report Bristol City Council; 2012.
- 10 Bornioli A, Bray I, Pilkington P, *et al*. The effectiveness of a 20 mph speed limit intervention on vehicle speeds in Bristol, UK: A non-randomised stepped wedge design. *J Transp Health* 2018;11:47–55.
- 11 Hemming K, Haines TP, Chilton PJ, *et al*. The stepped wedge cluster randomised trial: rationale, design, analysis, and reporting. *BMJ* 2015;350:h391.
- 12 Hussey MA, Hughes JP. Design and analysis of stepped wedge cluster randomized trials. *Contemp Clin Trials* 2007;28:182–91.
- 13 Hauer E. Observational before/after studies in road safety. estimating the effect of highway and traffic engineering measures on road safety. *Emerald Group Publishing Limited* 1997.
- 14 Department for Transport. Reported road casualties great Britain, annual report: 2017; 2017.