



Risk and protective factors associated with gang affiliation among high-risk youth: a public health approach

Dawn Delfin McDaniel

► An additional appendix is published online only. To view this file please visit the journal online (<http://injuryprevention.bmj.com/content/18/4.toc>).

Correspondence to

Dr Dawn D McDaniel, Epidemic Intelligence Service Officer assigned to the Division of Violence Prevention, National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, 4770 Buford Hwy NE, MS F64, Atlanta, GA 30341, USA; dawn.mcdaniel@cdc.hhs.gov

The findings and conclusions in this manuscript are those of the author and do not necessarily represent the views of the Centers for Disease Control and Prevention/the Agency for Toxic Substances and Disease Registry.

Accepted 8 October 2011
Published Online First
11 January 2012



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://injuryprevention.bmj.com/site/about/unlocked.xhtml>

ABSTRACT

Background Gang violence accounted for 20% of homicides in large cities from 2002 to 2006. Preventing gang affiliation (ie, youth who either desire or have gang membership) might reduce subsequent gang activity. Previous research has focused on identifying risk factors for gang affiliation; however, little information is available on protective factors.

Aim To identify risk and protective factors to provide more direction for gang violence prevention strategies.

Methods The author analysed cross-sectional survey data from 4131 youths in grades 7, 9, 11 and 12. Data were collected in 2004 from students in a high-risk, urban public school district. Regression analyses were conducted to assess the association between gang affiliation and alcohol and drug use, delinquency, depressed mood, suicidal ideation, peer victimisation, parental monitoring and positive reinforcement, adult, family and peer support, coping skills, and school connectedness. Analyses were controlled for sex, race/ethnicity and age.

Results An estimated 7% of youths were gang affiliated. Adjusting for all factors, gang affiliation was positively associated with engaging in any delinquent behaviours (prevalence OR: 2.07; 95% CI 1.18 to 3.64), frequent alcohol use (OR: 2.62; 95% CI 1.85 to 3.72) and frequent drug use (OR: 1.95; 95% CI 1.15 to 3.29). Gang affiliation was negatively associated with moderate levels of parental monitoring (OR: 0.67; 95% CI 0.54 to 0.85) and coping skills (OR: 0.54; 95% CI 0.42 to 0.71).

Conclusions The findings suggest the potential benefit of increasing parental monitoring and coping skills and reducing delinquency, alcohol use and drug use to prevent gang affiliation.

INTRODUCTION

From 2002 to 2006, gangs were responsible for approximately 20% of homicides in the 88 largest US cities,¹ and gang violence has been perceived as reaching 'epidemic' levels in many urban areas.² Furthermore, gang members are more likely than their non-gang-affiliated peers to engage in crime and violence,³⁻⁴ which increases their risk of violence-related injuries and death.⁵⁻⁶ While only an estimated 5% of the US population has ever joined a gang,⁷ gang membership has reached 14–30% of the population in many urban areas.⁸⁻⁹ The impact of gang membership on a youth's health is tremendous and preventing gang membership or the desire for gang affiliation early in adolescence might significantly reduce subsequent gang-related violence. This study explores a public health approach to understanding the factors that increase

risk for, as well as protect against, gang affiliation among youth who reside in a high-risk community (ie, a community with high levels of serious crime, poverty and unemployment).

Traditionally, initiatives designed to reduce gang affiliation have not focused on primary prevention strategies; that is, strategies aimed at preventing gang affiliation before gangs begin recruiting youth. These strategies often attempt to prevent factors that might increase risk of gang affiliation while promoting factors that might lower risk. Risk factors for gang affiliation include friendships with delinquent peers, lack of parental monitoring, negative life events (eg, loss of parent), and positive attitudes towards delinquency or engaging in delinquency (eg, violence perpetration).⁷⁻¹⁰ Other risky behaviours associated with gang affiliation include alcohol use and drug use.¹¹ To date, little research has focused on mental health related factors, such as depression and suicidal ideation;¹¹ these factors might be important since they are common among youth who engage in delinquency.¹² Several studies have also shown that the probability of gang affiliation increases as risk factors accumulate.¹⁰⁻¹¹

In contrast to the amount of literature on risk factors for gang affiliation, the literature on protective factors is limited. (The term 'protective factor' often refers to a factor that buffers the effects of risk. However, for this study, the term refers to factors found to be negatively associated with gang affiliation.) The few studies that have embarked on this topic have, in general, found that youth who have strong social skills, a supportive family (eg, parental monitoring, warmth and control) and connections with religion/religious institutions are less likely to seek gang affiliation than their peers¹³⁻¹⁵; however, more research is still needed to understand the protective factors for youth in high-risk communities, especially with regard to understudied factors, such as positive peer support and school commitment.¹⁶ This study assesses risk and protective factors associated with gang affiliation among a high-risk youth population to better inform primary prevention strategies.

METHODS

Study design

This study used data provided by the Centers for Disease Control and Prevention's 'Youth Violence Survey: Linkages among Different Forms of Violence' study.¹⁷ The Youth Violence Survey was a cross-sectional study of youth in a high-risk community that examined various types of

violence and their common risk and protective factors. These factors addressed the different domains of the social–ecological model, which includes individual, peer, family and community level factors.¹⁸ This survey also captured information on gang affiliation.

Study population

The Youth Violence Survey was administered to students in grades 7, 9, 11 and 12 in a high-risk community. The high-risk community was identified using multiple indicators of risk (ie, nationally the school district was among the highest 25 in poverty, the highest 15 in single-parent families, the highest 10 in serious crime rates and the highest 35 in rate of unemployment). This site was selected based on these demographic characteristics and pragmatic considerations (eg, commitment to study and feasibility of obtaining adequate census of students in targeted grades). While the site will not be named to protect its confidentiality, the community was racially and ethnically diverse with a population of less than 250 000. There were 16 public schools within this community's school district and all schools participated.

The study received Institutional Review Board approval from the Centers for Disease Control and Prevention and ORC Macro International. Prior to data collection, written parental consent and student assent were obtained from all students: 14% of students did not return the consent form and 1% of students refused to participate. Non-English speaking parents were offered consent forms in the major languages represented in the community, including Spanish.

Data collection for this study was undertaken in April 2004. Students voluntarily completed the anonymous self-administered 174-item questionnaire during one class period. Students received a \$5 gift card for returning the parental permission form (regardless of whether parental permission was granted) and another \$5 gift card after completing the survey. Those students who did not want to participate were assigned individual deskwork by their teacher. Among the 5098 students who met eligibility for the study, a total of 4131 (81%) chose to participate: 1491 students were in grade 7 (83% participation rate), 1117 students were in grade 9 (73.4% participation rate) and 1523 students were in grades 11 and 12 (79% participation rate). Further information about the study design, the population and the setting is available in the study by Swahn *et al.*¹⁷

Measures

Outcome variable

The outcome variable for this study, gang affiliation, was based on one item in which students were asked how they felt about joining a gang. The students were not given clarification regarding the definition of a gang. Responses were dichotomised to distinguish between gang-affiliated students who reported either the desire to join a gang or have an active gang membership with no intention of leaving the gang and non-gang-affiliated students who reported no desire to join or actively participate in a gang.

Main explanatory variables of interest

Thirteen variables were examined to identify their association with gang affiliation. These variables spanned the domains of individual, peer, family and community level characteristics. Six factors were assessed to determine whether they had a positive association (ie, act as potential risk factors). These factors included having a depressed mood in the past 30 days, having suicidal ideation in the past year, having consumed alcohol at

least once in the past year, having used drugs at least once in the past year, engaging in any delinquent behaviour in the past year and having been victimised in the past year. Seven variables were assessed to determine whether they had a negative association with the outcome variable (ie, act as potential protective factors). These variables included having the confidence to cope with conflict, having peer support, having parents who have used positive reinforcement in the past 30 days, having parents who have used monitoring strategies in the past 30 days, having family support, having adult support at school and feeling connected to school.

Details of how these variables were coded from the survey are shown in the online supplementary appendix A. In general, these variables exhibited high internal consistency (see Logan *et al* for further description¹⁹). Multi-item variables were re-classified to account for the distribution of the data in a conservative manner while retaining variability when possible.

Statistical analysis

Basic descriptive analyses were conducted to characterise non-gang- and gang-affiliated students. Logistic regression was used to assess crude associations between the outcome variable and demographic variables as well as the main variables of interest. Associations were reported via prevalence ORs. Robust variance estimates were used to account for potential effects from cluster correlation within each school. Multiple logistic regression was used to determine the adjusted ORs for each of the main variables of interest. This model adjusted for demographic factors (ie, age, sex and race/ethnicity) to control for potential developmental and sociocultural influences.²⁰ Since youths can have more than one risk or protective factor, we also adjusted for all of these factors in the model to provide more accurate individual associations between the factors and the outcome variable. Furthermore, we assessed for collinearity using variable inflation factor scores. Variable inflation factor scores fell within acceptable ranges (<2), which suggests that collinearity among the variables used in the models was not an issue.

The proposed risk and protective factors were summed to create cumulative indices. For these indices, all variables were rendered dichotomous (see online supplementary appendix A for more details). Multiple logistic regressions were used to determine the association of these indices with gang affiliation, controlling for age, sex and race/ethnicity. Also, the cumulative risk index was dichotomised into a risk grouping variable by a median split to distinguish those at high risk (≥ 3 risk factors) from those at low risk (≤ 2 risk factors) for subsequent analyses. This variable was used in a model that included interaction terms between this risk grouping variable and the salient protective factors from the adjusted model.

To illustrate the potential benefit of having the salient protective factors at high and low levels of risk, another multiple logistic regression was run including the risk grouping variable and key protective factors. The estimated predicted probabilities for gang affiliation were determined for youth at high and low risk with no protective factors and with each combination of protective factors. Fisher's exact tests with a Bonferroni correction for multiple comparisons were used to compare these probabilities.

RESULTS

Descriptive characteristics of the study population are reported in table 1. The majority of this youth population was female, of ethnic minority status and of high school age (ie, 14–18 years or older). An estimated 7.1% of the youths were gang

Table 1 Frequencies and crude ORs for demographic, risk and protective factors for gang affiliation (N=4131)

Variable	Total N (%)	Gang-affiliated students (N=294)	
		N (%)	Crude OR (95% CI)
Demographic factors			
Age			
12–13 years	1222 (29.58)	95 (7.77)	Referent
14–15 years	1180 (28.56)	105 (8.90)	1.16 (0.72 to 1.88)
16 years or older	1705 (41.27)	92 (5.40)	0.68 (0.38 to 1.22)
Missing	24 (0.58)	2 (8.33)	1.08 (0.25 to 4.62)
Sex			
Female	2127 (51.49)	113 (5.31)	Referent
Male	1982 (47.98)	179 (9.03)	1.77 (1.37 to 2.28)
Missing	22 (0.53)	2 (9.09)	1.78 (0.38 to 8.47)
Race/ethnicity			
White, non-Hispanic	922 (22.32)	43 (4.66)	Referent
African American, non-Hispanic	935 (22.63)	88 (9.41)	2.12 (1.21 to 3.73)
Hispanic	1809 (43.79)	128 (7.08)	1.56 (1.05 to 2.31)
Others, non-Hispanic	416 (10.07)	33 (7.93)	1.76 (1.08 to 2.86)
Missing	49 (1.19)	2 (4.08)	0.87 (0.21 to 3.57)
Potential risk factors			
Depressed mood†			
No	2464 (59.65)	154 (6.25)	Referent
Yes	1667 (40.35)	140 (8.40)	1.43 (1.07 to 1.92)
Suicidal ideation			
No	3416 (82.69)	210 (6.15)	Referent
Yes	715 (17.31)	84 (11.75)	2.03 (1.62 to 2.55)
Alcohol use			
Never	2009 (48.63)	81 (4.03)	Referent
Once a year to once a month	1273 (30.82)	83 (6.52)	1.66 (1.11 to 2.47)
2–3 days a month	328 (7.94)	35 (10.67)	2.84 (1.76 to 4.58)
1–2 days a week or more	391 (9.47)	84 (21.48)	6.51 (4.64 to 9.15)
Missing	130 (3.15)	11 (8.46)	2.20 (0.88 to 5.52)
Drug use			
Never	3090 (74.80)	139 (4.50)	Referent
Once a year to once a month	409 (9.90)	41 (10.02)	2.37 (1.50 to 3.74)
2–3 days a month	130 (3.15)	17 (13.08)	3.19 (1.84 to 5.56)
1–2 days a week or more	409 (9.90)	84 (20.54)	5.49 (3.68 to 8.17)
Missing	93 (2.25)	13 (13.98)	3.45 (1.79 to 6.64)
Delinquency			
Never	1503 (36.38)	29 (1.93)	Referent
1–3 times	1478 (35.78)	80 (5.41)	2.91 (1.53 to 5.52)
4–9 times	447 (10.82)	56 (12.53)	7.28 (4.38 to 12.10)
10 or more times	645 (15.61)	127 (19.69)	12.46 (7.38 to 21.05)
Missing	58 (1.40)	2 (3.48)	1.82 (0.40 to 8.17)
Peer victimisation			
No	1880 (45.51)	114 (7.45)	Referent
Yes	2251 (54.49)	180 (8.00)	1.35 (1.04 to 1.74)
Potential protective factors			
Coping skills			
Low level of confidence	1475 (35.71)	197 (13.36)	Referent
Moderate level of confidence	1472 (35.63)	75 (5.10)	0.35 (0.27 to 0.44)
High level of confidence	1184 (28.66)	22 (1.86)	0.12 (0.08 to 0.19)
Peer support			
No	2050 (49.62)	173 (8.44)	Referent
Yes	2081 (50.38)	121 (5.81)	0.67 (0.52 to 0.86)

Continued

Table 1 Continued

Variable	Total N (%)	Gang-affiliated students (N=294)	
		N (%)	Crude OR (95% CI)
Parental positive reinforcement			
No	1527 (36.96)	130 (8.51)	Referent
Yes	2604 (63.04)	164 (6.30)	0.72 (0.56 to 0.93)
Parental monitoring			
Low level	1480 (35.83)	175 (11.82)	Referent
Moderate level	1332 (32.24)	75 (5.63)	0.44 (0.36 to 0.55)
High level	1319 (31.93)	44 (3.34)	0.26 (0.20 to 0.33)
Family support			
No	1823 (44.13)	173 (9.49)	Referent
Yes	2308 (55.87)	121 (5.24)	0.53 (0.41 to 0.68)
Adult support at school			
No	268 (6.49)	33 (12.31)	Referent
Yes	3863 (93.51)	261 (6.76)	0.52 (0.28 to 0.96)
School connectedness			
No	867 (20.99)	75 (8.65)	Referent
Yes	3264 (79.01)	219 (6.71)	0.76 (0.62 to 0.92)

†The missing response category was added to the referent group for all dichotomous variables and variables divided by tertiles.

affiliated: 2.4% of the youths reported the desire to join a gang and 4.8% of the youths reported active gang membership with no intention of leaving the gang. A majority also reported having consumed alcohol in the past 12 months (48.2%), having participated in delinquent behaviour in the past 12 months (62.2%) and having been victimised by a peer in the past 12 months (54.5%). However, a large proportion of the study population also reported having at least a moderate level of confidence to cope with conflict (64.3%), having parents who have provided positive reinforcement in the past 30 days (63%), having parents who monitored their activities at least at a moderate level in the past 30 days (64.2%), having adult support at school (93.5%) and feeling connected to school (79%).

The crude associations between the predictor variables and the outcome variable in table 1 revealed that gang affiliation was most common among youth of ethnic minority status, males, those who had a depressed mood, reported suicidal ideation, used alcohol or drugs, were delinquent, and had experienced peer victimisation. Gang affiliation was least common among youth who had confidence in their coping skills, peer support, parental positive reinforcement, parental monitoring, family support, adult support at school and school connectedness.

The adjusted ORs for all predictor variables are presented in table 2. Alcohol use that occurs at least 2–3 days a month, drug use that occurs at least 1–2 days a week and any delinquency remained positively associated with gang affiliation. Having at least a moderate level of confidence with regard to coping with conflict and having at least a moderate level of parental monitoring remained negatively associated with gang affiliation.

The findings on cumulative risk and protective factor indices are presented in table 3. Over half of the study population reported two or more risk factors. Those youths with four or more risk factors were nearly six times more likely to be gang affiliated than those with zero to one risk factor. Also, those with three or less protective factors were 5.6 times as likely to be gang affiliated than those who had four or more protective factors reported. The cumulative risk and protective factor indices were analysed in an adjusted model, controlling for demographic characteristics. The odds of gang affiliation were

Table 2 Adjusted ORs and 95% CIs for demographic, risk and protective factors for gang affiliation (N=4131)

Variable	Adjusted OR (95% CI)
Demographic factors	
Age†	
12–13 years	Referent
14–15 years	0.89 (0.59 to 1.34)
16 years or older	0.45 (0.29 to 0.71)
Sex	
Female	Referent
Male	1.35 (1.00 to 1.82)
Race/ethnicity	
White, non-Hispanic	Referent
African American, non-Hispanic	2.37 (1.51 to 3.70)
Hispanic	1.45 (1.09 to 1.91)
Others, non-Hispanic	1.79 (1.14 to 2.80)
Potential risk factors	
Depressed mood	
No	Referent
Yes	1.20 (0.87 to 1.65)
Suicidal ideation	
No	Referent
Yes	1.39 (0.93 to 2.08)
Alcohol use	
Never	Referent
Once a year to once a month	1.15 (0.80 to 1.64)
2–3 days a month	1.48 (1.04 to 2.13)
1–2 days a week or more	2.62 (1.85 to 3.72)
Drug use	
Never	Referent
Once a year to once a month	1.57 (0.98 to 2.51)
2–3 days a month	1.67 (0.94 to 2.97)
1–2 days a week or more	1.95 (1.15 to 3.29)
Delinquency	
Never	Referent
1–3 times	2.07 (1.18 to 3.64)
4–9 times	3.36 (1.88 to 6.02)
10 or more times	5.10 (3.20 to 8.11)
Peer victimisation	
No	Referent
Yes	1.07 (0.81 to 1.41)
Potential protective factors	
Coping skills	
Low level of confidence	Referent
Moderate level of confidence	0.54 (0.42 to 0.71)
High level of confidence	0.30 (0.19 to 0.47)
Peer support	
No	Referent
Yes	1.03 (0.75 to 1.40)
Parental positive reinforcement	
No	Referent
Yes	1.08 (0.87 to 1.34)
Parental monitoring	
Low level	Referent
Moderate level	0.67 (0.54 to 0.85)
High level	0.61 (0.43 to 0.86)
Family support	
No	Referent
Yes	0.82 (0.57 to 1.17)

Continued

significantly higher as a youth accumulated more than two risk factors; however, as the youth acquired over five protective factors the odds of being affiliated with a gang was significantly lowered. No significant associations were found for the inter-

Table 2 Continued

Variable	Adjusted OR (95% CI)
Adult support at school	
No	Referent
Yes	0.74 (0.40 to 1.37)
School connectedness	
No	Referent
Yes	1.15 (0.91 to 1.46)

The model was adjusted for all demographic factors, potential risk factors and potential protective factors.

†The missing response category was added to the referent group for all variables.

action terms between risk level and the significant protective factors, suggesting that these protective factors did not buffer risk. Instead, the protective factors had a similar negative association with gang affiliation for all youth regardless of risk level.

The predicted probabilities for gang affiliation based on various combinations of the significant protective factors (ie, confidence in coping skills and parental monitoring) and risk level are presented in table 4. Youth who reported having either coping skills alone or coping skills in combination with parental monitoring had a significantly lower probability for gang affiliation than those youth with neither protective factor, regardless of risk level ($p < 0.01$). Also, youth who were considered high risk with both protective factors had significantly lower probabilities for gang affiliation than youth at low risk who did not have either protective factor ($p < 0.05$). Youth at high risk with coping skills alone had similar probabilities for gang affiliation as youth at low risk with neither protective factor.

DISCUSSION

The finding that gang affiliation is associated with engaging in delinquent activities, particularly illicit alcohol and drug use, is one of the most consistent findings in the gang literature.^{4 11} While the temporal relationship of these variables cannot be explained using the current study design, prior research using longitudinal data has shown that delinquency and illicit substance use have an 'enhancing' relationship with gang affiliation.¹¹ For example, it has been found that early involvement in these antisocial behaviours elevates the risk for subsequent gang

Table 3 Frequencies and adjusted ORs for cumulative risk and protective factors for gang affiliation (N=4131)

Variable	Total N (%)	Gang-affiliated students (N=294)		
		N (%)	Adjusted OR	95% CI
Cumulative risk index				
0	532 (12.88)	6 (1.13)	Referent	
1	1002 (24.26)	33 (3.29)	2.63	0.99 to 7.00
2	1075 (26.02)	59 (5.49)	4.37	1.64 to 11.60
3	792 (19.17)	81 (10.22)	9.31	3.53 to 24.58
4	493 (11.93)	70 (14.20)	14.81	5.62 to 39.02
5–6	237 (5.73)	45 (18.99)	21.51	7.89 to 58.64
Cumulative protective index				
0–3	481 (11.64)	125 (25.99)	Referent	
4	727 (17.60)	69 (9.50)	0.81	0.63 to 1.05
5	877 (21.23)	57 (6.50)	0.63	0.49 to 0.81
6	793 (19.20)	32 (4.04)	0.44	0.31 to 0.61
7	705 (17.07)	11 (1.56)	0.21	0.12 to 0.36

Age, sex and race/ethnicity were controlled for in the models.

Table 4 Predicted probabilities of gang affiliation for youth with various combinations of the significant protective factors and risk level

Protective factors	Low risk	High risk
Neither parental monitoring nor coping skills	^A 0.10	^B 0.26
Parental monitoring alone	^C 0.05	^D 0.15
Coping skills alone	^E 0.03	^F 0.10
Both parental monitoring and coping skills	^G 0.02	^H 0.05

Comparisons of probabilities		
Low risk	High risk	High risk × Low risk
C×A	D×B**	H×A*
E×A**	F×B**	
G×A**	H×B**	

Low risk=0–2 risk factors. High risk=3–6 risk factors. To compare the predicted probabilities, Fisher's exact tests were used with a Bonferroni correction for multiple comparisons.

* $p < 0.05$, ** $p < 0.01$.

affiliation by reducing the possibility for positive social connections and increasing opportunities for delinquent peer associations. Furthermore, the reciprocal relationship is also thought to exist where gang affiliation 'facilitates' an increase in these antisocial behaviours (eg, excessive substance abuse, more severe criminal behaviour) through peer pressure.¹¹ Reducing the risk of initiating illicit substance use or delinquency early in preadolescence or adolescence might disrupt the initial trajectory towards gang affiliation or even potentially break current gang-related connections that could further escalate these behaviours.

This study also showed the potential benefit of emphasising protective factors in prevention efforts. Two protective factors, youth's confidence in their coping skills and parental monitoring, were negatively associated with gang affiliation. Youths at high risk who had the two key protective factors had lower probabilities for gang affiliation than youths with neither of these factors at low risk. Also, youths at high risk with only one protective factor, confidence in their coping skills, had similar probabilities for gang affiliation as youths at low risk with neither factor. For youths in high-risk contexts, a focus on reducing risk factors may not be feasible in all situations due to the number of risk exposures and the difficulty in changing certain risk factors. Therefore, if it is not possible to reduce risk, adding one or two key protective factors may make youth at high risk appear similar to those at low risk. Other studies have documented a similar value in the use of protective factors against other forms of violence.^{21–23}

Limitations

This study was subject to several limitations. First, the cross-sectional nature of these data limited our ability to understand the causal ordering of the risk and protective factors and the outcome, gang affiliation. For instance, we were unable to determine whether confidence in coping with conflict prevented gang affiliation or whether youth developed poor coping strategies while affiliated with gangs. We also did not attempt to disentangle the complex relationships between gang affiliation, delinquency, alcohol use and drug use. Future prospective research would provide a better understanding of the relationship of these correlates. Second, surveying gang-affiliated youth in school may introduce a selection bias, and a non-response bias analysis was not done in this study. Further work that assesses whether these findings can be replicated in samples of youth who have dropped out of school, are truant or are adjudicated is important. Third, the survey did not define gang affiliation for respondents; therefore, gang affiliation in this study is likely to

have a broad meaning. Past research has shown that using a more restrictive definition leads to more precision in distinguishing gangs from other youth groups.⁷ Finally, this study is limited to a high-risk population and the findings cannot, by this study design, be generalised to other populations.

Prevention strategies

This study has several important implications for prevention efforts. Since youth with both parental monitoring and coping skills had the lowest probability of gang affiliation, programmes that offer parent training that are focused on skills relevant to effectively monitoring children, in addition to helping youth develop strategies to cope with conflict, may be most beneficial for preventing gang affiliation. Several primary prevention programmes aimed at reducing delinquent behaviour have shown positive effects.^{24–25} Examples of evidence-based prevention programmes that address coping skills (ie, interpersonal skills, such as talking out disagreements, and intrapersonal skills, such as controlling anger) include the Promoting Alternative THinking Strategies (PATHS) curriculum and I Can Problem Solve.^{24–25} Examples of concurrent parent and child programmes targeting both coping skills and effective parenting include Strengthening Families Program, Fast Track, Prevention Treatment Program and Incredible Years.²⁵ Of these programmes, only the Prevention Treatment Program showed reductions in gang affiliation.²⁶ While the other programmes have not evaluated their impact on gang affiliation, given the shared aetiology of delinquency, drug and substance use, and gang affiliation, these programmes would likely be promising.

Future directions

Using a public health approach, we identified the potential of increasing a youth's confidence in their ability to cope with conflict and parental monitoring and reducing their delinquency, alcohol use and drug use to prevent gang affiliation. While studies that identify risk and protective factors represent one of the initial steps of the public health model, further work needs to be done to develop and evaluate primary prevention

What is already known on the subject

- ▶ Risk factors for gang affiliation include friendship with delinquent peers, lack of parental monitoring, negative life events and positive attitudes towards delinquency or engaging in delinquency.
- ▶ Other risky behaviours associated with gang affiliation include alcohol use and drug use.
- ▶ The probability of gang affiliation increases as risk factors accumulate.

What this study adds

- ▶ This study showed the potential value of emphasising protective factors when assessing factors associated with gang affiliation.
- ▶ The findings suggest the potential benefit of increasing parental monitoring and coping skills and reducing delinquency, alcohol use and drug use to prevent gang affiliation.

programmes that help reduce gang affiliation. Also, more work is needed to help disseminate programmes that can potentially prevent gang affiliation. Dissemination may be best accomplished with the inclusion of public health professionals in the planning and implementation of such programmes. These public health professionals may be able to assist in the necessary on-going collaboration with other sectors that traditionally work with gang-affiliated youth, such as criminal justice, to be most strategic about ways of combining community efforts to reduce gang affiliation.

Acknowledgements I wish to thank Dr J Logan for his contributions to this research.

Funding This study was supported by Centers for Disease Control and Prevention.

Competing interests None.

Ethics approval Ethics approval was provided by Centers for Disease Control and Prevention (CDC) and ORC Macro International.

Contributors The Youth Violence Survey was developed by the CDC and was implemented in 2004 by contractors at ORC Macro International.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data were obtained from the CDC's Youth Violence Survey.

REFERENCES

1. **Pyrooz D.** Structural covariates of gang homicide in large U.S. cities. *J Res Crime Delinquen* 2011;**48**:1–30.
2. **Hutson H,** Anglin D, Kyriacou D, *et al.* The epidemic of gang-related homicides in Los Angeles County from 1979 through 1994. *JAMA* 1995;**274**:1031–6.
3. **Thornberry TP,** Lizotte AJ, Krohn MD, *et al.* Causes and consequences of delinquency: findings from the Rochester Youth Development Study. In: Thornberry TP, Krohn MD, eds. *Taking Stock of Delinquency*. New York: Kluwer Academic/Plenum Publishers, 2003:11–46.
4. **Gatti U,** Tremblay RE, Vitaro F, *et al.* Youth gangs, delinquency and drug use: a test of the selection, facilitation, and enhancement hypotheses. *J Child Psychol Psychiatry* 2005;**46**:1178–90.
5. **Taylor T,** Peterson D, Esbensen F, *et al.* Gang membership as a risk factor for adolescent violent victimization. *J Res Crime Delinquen* 2007;**44**:351–80.
6. **Spano R,** Freilich J, Bolland J. Gang membership, gun carrying, and employment: applying routine activities theory to explain violent victimization among inner city, minority youth living in extreme poverty. *Justice Quarterly* 2008;**25**:381–410.
7. **Klein M,** Maxson C. *Street Gang Patterns and Policies*. New York: Oxford University Press, 2006.
8. **Thornberry T.** Membership in youth gangs and involvement in serious and violent offending. In: Loeber R, Farrington D, eds. *Serious and Violent Offenders: Risk Factors and Successful Intervention*. Thousand Oaks: Sage, 1998:147–66.
9. **Thornberry T,** Huizinga D, Loeber R. The causes and correlates studies: findings and policy implications. *Juvenile Justice Bulletin* 2004;**9**:3–19.
10. **Hill K,** Howell J, Hawkins J, *et al.* Childhood risk factors for adolescent gang membership: results from the Seattle social development project. *J Res Crime Delinquen* 1999;**36**:300–22.
11. **Thornberry T.** *Gangs and Delinquency in Developmental Perspective*. Cambridge: Cambridge University Press, 2003.
12. **Thompson MP,** Ho CH, Kingree JB. Prospective associations between delinquency and suicidal behaviors in a nationally representative sample. *J Adolesc Health* 2007;**40**:232–7.
13. **Li X,** Stanton B, Pack R, *et al.* Risk and protective factors associated with gang involvement among urban African American adolescents. *Youth Soc* 2002;**34**:172–94.
14. **Katz C,** Fox A. Risk and protective factors associated with gang-involved youth in Trinidad and Tobago. *Revista Panamericana de Salud Pública* 2010;**27**:187–202.
15. **Maxson C,** Whitlock M, Klein M. Vulnerability to street gang membership: implications for practice. *Soc Serv Rev* 1998;**72**:70–91.
16. **Sharkey J,** Shekhtmeyster Z, Chavez-Lopez L, *et al.* The protective influence of gangs: can schools compensate? *Aggression and Violent Behavior* 2011;**16**:45–54.
17. **Swahn MH,** Simon TR, Arias I, *et al.* Measuring sex differences in violence victimization and perpetration within date and same-sex peer relationships. *J Interpers Violence* 2008;**23**:1120–38.
18. **Brofenbrenner U.** *The Ecology of Human Development*. Cambridge: Harvard University Press, 1979.
19. **Logan J,** Crosby A, Hamburger M. Suicidal ideation, friendships with delinquents, social and parental connectedness, and differential associations by sex: findings among a high-risk pre/early adolescent population. *Crisis*. Published Online First: 6 July 2011. doi:10.1027/0227-5910/a000091.
20. **Esbensen F,** Peterson D, Taylor T, *et al.* *Youth Violence: Sex and Race Differences in Offending, Victimization, and Gang Membership*. Philadelphia: Temple University Press, 2010.
21. **Borowsky IW,** Ireland M, Resnick MD. Adolescent suicide attempts: risk and protectors. *Pediatrics* 2001;**107**:485–93.
22. **Borowsky IW,** Hogan M, Ireland M. Adolescent sexual aggression: risk and protective factors. *Pediatrics* 1997;**100**:E7.
23. **Borowsky IW,** Resnick MD, Ireland M, *et al.* Suicide attempts among American Indian and Alaska Native youth: risk and protective factors. *Arch Pediatr Adolesc Med* 1999;**153**:573–80.
24. **Hahn R,** Fuqua-Whitley D, Wethington H, *et al.* The effectiveness of universal school-based programs for the prevention of violent and aggressive behavior: a report on recommendations of the Task Force on Community Preventive Services. *MMWR Recomm Rep* 2007;**56**:1–12.
25. **Elliott DS.** *Blueprints for Violence Prevention (Vols. 1–11)*. Boulder: Center for the Study and Prevention of Violence, Institute of Behavioral Science, University of Colorado, 1997.
26. **McCord J,** Tremblay RE, Vitaro F, *et al.* Boys' disruptive behaviour, school adjustment, and delinquency: the Montreal prevention experiment. *Int J Behav Dev* 1994;**17**:739–52.

Still undecided about using a cellphone while driving?

For the past 4 years, California has banned drivers from using handheld cellphones. A study reveals that traffic deaths involving cellphones had fallen by 47%. Deaths among drivers using hands-free phones dropped at a similar rate. The study examined deaths in the 2 years before and 2 years after the cellphone ban took effect, and found a similar drop in injuries. The explanation for the exceptional effectiveness of this law is that it was enforced: there were 460 487 convictions in 2011 up 52% from 2009. The Senator who proposed the law wants to increase the penalty from \$20 to \$30 for a first offense, but with court and other fees, the net cost, currently, is at least \$159.

APPENDIX A. Description of Variables

(Web Only File)

Domain	Explanation
<i>Individual Level</i>	
Coping skills	<p>This variable assessed a youth’s average level of confidence in his or her ability to use various coping strategies (i.e., stay out of fights by choosing other solutions, talk out a disagreement, calm down when mad, ignore someone’s teasing, avoid a fight by walking away, apologize to the other students, and seek help from an adult). These items had high internal consistency (Cronbach’s alpha = 0.88). The responses for each item were coded on a scale from 0 to 2 (i.e., 0 = “no confidence;” 1 = “somewhat confident;” 2 = “very confident”). The mean scale score was calculated for each youth. Based on the tertiles of the distribution of these values, youth were placed in one of three categories: low level of coping, moderate level of coping, high level of coping.</p>
Depressed mood	<p>This variable was dichotomized to capture any symptoms of a depressed mood. Youths were asked how many times they had been sad, grouchy or irritable or moody, and hopeless about the future in the past 30 days (Cronbach’s alpha = 0.77). To capture a youth having any of these symptoms at a clinically relevant level, each respondent was categorized based on his or her most severe response. Therefore, youths reporting “often” or “always” to any of the symptoms were categorized in the depressed mood group.</p>
Suicidal ideation	<p>This variable was comprised of one item, “Did you ever seriously consider attempting suicide?”, which was used to assess suicidal ideation in the past 12 months. Response options were “yes” and “no”.</p>

Alcohol use	This variable used one item to capture the frequency youth had at least one drink of alcohol in the past 12 months. Response options were coded on a scale of 1 to 4 (i.e., 1 = “never;” 2 = “once a month or less;” 2 = “2-3 days a month;” 3 = “1-2 days a week or more”).
Drug use	This variable used one item to capture the frequency youth had used inhalants (glue or solvents) or illegal drugs, such as marijuana, cocaine, and heroin in the past 12 months. Response options were coded on a scale of 1 to 4 (i.e., 1 = “never;” 2 = “once a month or less;” 2 = “2-3 days a month;” 3 = “1-2 days a week or more”).
Delinquency	Youth were considered delinquent if they reported engaging in at least one of the following behaviors: deliberately damage property that did not belong to them, hurt someone badly enough to need bandages or care from a doctor or nurse, steal things, use or threatened to use a weapon to get something from someone, and sell marijuana or other drugs. These items had high internal consistency (Cronbach’s alpha = 0.75). Response options ranged from 1 to 4 (1 = “never;” 2 = “1 or 2 times;” 3 = “3 or 4 times;” 4 = “5 or more times”). To capture the frequency of these behaviors, each respondent was categorized based on his or her most severe response.
Peer victimization	This variable was dichotomized to distinguish between those who have never been victimized and those who have had a peer do at least one of the following to them in the past 12 months: damage something that belonged to them, say things to hurt their feelings on purpose, threatened to hit or throw something at them, insulted them in front of others, scratched them, put down their looks, hit or slapped them, slammed them or held them against a wall, kicked them, pushed, grabbed, or shoved them, forced them to have sex or to do something sexual that they did not want to do, threw something at them that could hurt, punched or hit them with something that could hurt, threatened or injured them with a knife or gun, and hurt them badly enough to need bandages or care from a doctor or nurse. These items had high internal consistency (Cronbach’s alpha = 0.92).

<i>Peer level</i>	
Peer support	This variable was dichotomized. Youths were considered to have peer support if they agreed “a lot” to any of the following items: “I have friends I can talk to, who care about my feelings and what happens to me; “I have friends I can talk to, who give good suggestions and advice about my problems;” and “I have friends who help me with practical problems, like how to get somewhere, or help me with a job or project” (Cronbach’s alpha = 0.90).
<i>Family level</i>	
Parental positive reinforcement	This variable was dichotomized. Youths were considered to have consistent parental positive reinforcement in the past 30 days if they reported that their parents “almost always” did any of the following to show approval: say something nice, praise or approval; hug, pat on the back, or kiss; give rewards; give special privileges; or do a special activity (Cronbach’s alpha = 0.79).
Parental monitoring	This variable captured the consistency of parent’s use of monitoring in the past 30 days. Parental monitoring strategies included: set curfew on school nights, set curfew on weekend, tell parents/guardian what you were doing when you were outside of the house, leave a note or call parents to let them know where you were going when they are not home, when you were out, tell parents who you were with when you were not at home, tell parents what you were doing when they are home (Cronbach’s alpha = 0.76). Response options ranged from 0 to 2 (0 = “No or almost never;” 1= “Sometimes;” 2=“Almost always”). The mean scale score was calculated. Based on the tertiles of the distribution of these values, youth were placed in one of three categories: low level of parental monitoring, moderate level of parental monitoring, high level of parental monitoring.

Family support	This variable was dichotomized. Youths were considered to be supported by their family if they agreed “a lot” with any of the following items: “There are people in my family I can talk to, who care about my feelings and what happens to me;” “There are people in my family I can talk to, who give good suggestions and advice about my problems;” and “There are people in my family who help me with practical problems, like helping me get somewhere or help me with a job or project” (Cronbach’s alpha = 0.91).
<i>Community level</i>	
Adult support at school	This variable was dichotomized. Youths were considered to be supported by their family if they reported agreeing “a lot” with any of the following items: “At school, there are adults I can talk to, who care about my feelings and what happens to me;” “At school, there are adults I can talk to, who give good suggestions and advice about my problems;” and “At school, there are adults who help me with practical problems, like helping me get somewhere, or helping me with a job or project” (Cronbach’s alpha=0.85).
School connectedness	This variable was dichotomized. Youths were considered to be connected to their school if they reported agreeing “a lot” with any of the following items: “I feel close to people at school;” “I feel part of the school;” “I feel happy to be at school” (Cronbach’s alpha=0.72).
Cumulative risk index	The risk variables were summed to create the index of cumulative risk factors. Non-dichotomous risk factors were dichotomized based on referent. The number of risk factors ranged from 0 to 6.
Cumulative protective index	The protective variables were summed to create the index of cumulative protective factors. Non-dichotomous protective factors were dichotomized based on referent. The number of protective factors ranged from 0 to 7.

Risk groups	This variable was created by splitting the cumulative risk index into two groups by its median: low risk group (0 to 2 risk factors) and high risk group (3 to 6 risk factors).
-------------	---